arXiv:1608.00373v1 [math.CO] 1 Aug 2016

The spectral excess theorem for graphs with few eigenvalues whose distance-2 or distance-1-or-2 graph is strongly regular

C. Dalfó[a], M.A. Fiol[a,b], J. Koolen[c]

[a]Universitat Politècnica de Catalunya, BarcelonaTech
Dept. de Matemàtiques, Barcelona, Catalonia
[b] Barcelona Graduate School of Mathematics, Catalonia
{cristina.dalfo,miguel.angel.fiol}@upc.edu
[c]University of Science and Technology of China
School of Mathematical Sciences
Hefei, Anhui, China
koolen@ustc.edu.cn

August 2, 2016

Abstract

We study regular graphs whose distance-2 graph or distance-1-or-2 graph is strongly regular. We provide a characterization of such graphs Γ (among regular graphs with few distinct eigenvalues) in terms of the spectrum and the mean number of vertices at maximal distance d from every vertex, where $d+1$ is the number of different eigenvalues of Γ. This can be seen as a another version of the so-called spectral excess theorem, which characterizes in a similar way those regular graphs that are distance-regular.

Keywords: Distance-regular graph; distance-2 graph; spectrum; predistance polynomials.

AMS subject classifications: 05C50, 05E30.

1 Preliminaries

Let Γ be a distance-regular graph with adjacency matrix $\boldsymbol{A}$ and $d+1$ distinct eigenvalues. In the recent works of Brouwer and Fiol [4, 15], it was studied the situation in which the distance-d graph Γ_d of Γ (or the Kneser graph K of Γ) with adjacency matrix $\boldsymbol{A}_d(=p_d(\boldsymbol{A}))$, where p_d is the distance-d polynomial, has fewer distinct eigenvalues than Γ. Examples are the so-called half antipodal (K with only one negative eigenvalue, up to multiplicity),

and antipodal distance-regular graphs (where K consists of disjoint copies of a complete graph).

Here we study the cases in which Γ has few eigenvalues and its distance-2 graph Γ_2 or its distance-1-or-2 graph $\Gamma_{1,2}$ are strongly regular. The main result of this paper is a characterization of such (partially) distance-regular graphs, among regular graphs with $d \in \{3, 4\}$ distinct nontrivial eigenvalues, in terms of the spectrum and the mean number of vertices at maximal distance d from every vertex. This can be seen as another version of the so-called spectral excess theorem. Other related characterizations of some of these cases were given by Fiol in [11, 12, 13]. For background on distance-regular graphs and strongly regular graphs, we refer the reader to Brouwer, Cohen, and Neumaier [3], Brouwer and Haemers [5], and Van Dam, Koolen and Tanaka [9].

Let Γ be a regular (connected) graph with degree k, n vertices, and spectrum $\operatorname{sp}\Gamma = \{\theta_0^{m_0}, \theta_1^{m_1}, \dots, \theta_d^{m_d}\}$, where $\theta_0 (= k) > \theta_1 > \cdots > \theta_d$, and $m_0 = 1$. In this work, we use the following scalar product on the $(d+1)$-dimensional vector space of real polynomials modulo $m(x) = \prod_{i=0}^{d}(x - \theta_i)$, that is, the minimal polynomial of $\boldsymbol{A}$.

$$\langle p, q\rangle_\Gamma = \frac{1}{n}\operatorname{tr}(p(\boldsymbol{A})q(\boldsymbol{A})) = \frac{1}{n}\sum_{i=0}^{d} m_i p(\theta_i)q(\theta_i), \qquad p, q \in \mathbb{R}_d[x]/(m(x)). \tag{1}$$

This is a special case of the inner product of symmetric $n \times n$ real matrices $\boldsymbol{M}$ and $\boldsymbol{N}$, defined by $\langle \boldsymbol{M}, \boldsymbol{N}\rangle = \frac{1}{n}\operatorname{tr}(\boldsymbol{M}\boldsymbol{N})$. The *predistance polynomials* $p_0, p_1, \dots, p_d$, introduced by Fiol and Garriga [17], are a sequence of orthogonal polynomials with respect to the inner product (1), normalized in such a way that $\|p_i\|_\Gamma^2 = p_i(k)$ (this makes sense since it is known that $p_i(k) > 0$ for any $i = 0, \dots, d$, see for instance [21]). As every sequence of orthogonal polynomials, the predistance polynomials satisfy a three-term recurrence of the form

$$xp_i = \beta_{i-1}p_{i-1} + \alpha_i p_i + \gamma_{i+1}p_{i+1} \qquad (i = 0, 1, \dots, d),$$

where the constants β_{i-1}, α_i, and γ_{i+1} are the Fourier coefficients of xp_i in terms of p_{i-1}, p_i, and p_{i+1}, respectively (and $\beta_{-1} = \gamma_{d+1} = 0$), beginning with $p_0 = 1$ and $p_1 = x$.

Some basic properties of the predistance polynomials and preintersection numbers are included in the following result (see Cámara, Fàbrega, Fiol, and Garriga [6], and Diego, Fàbrega, and Fiol [10]).

Lemma 1. *Let G be a k-regular graph with $d+1$ distinct eigenvalues and predistance polynomials $p_0, \dots, p_d$. Given an integer $\ell \geq 0$, let $\overline{\mathcal{C}}_\ell$ be the average number of circuits of length ℓ rooted at every vertex, that is, $\overline{\mathcal{C}}_\ell = \frac{1}{n}\sum_{i=0}^{d} m_i\theta_i^\ell$. Then,*

(*i*) *$p_0(x) = 1$, $p_1(x) = x$, $p_2(x) = \frac{1}{\gamma_2}(x^2 - \alpha_1 x - k)$.*

(*ii*) *For $i = 0, \dots, d$, the two highest terms of the predistance polynomial p_i are given by*

$$p_i(x) = \tfrac{1}{\gamma_1\cdots\gamma_i}[x^i - (\alpha_1 + \cdots + \alpha_{i-1})x^{i-1} + \cdots].$$

(*iii*) $\alpha_i + \beta_i + \gamma_i = k$, *for* $i = 0, \ldots, d$.

(*iv*) $\alpha_0 = 0$, $\beta_0 = k$, $\gamma_1 = 1$, $\alpha_1 = \overline{\mathcal{C}}_3/\overline{\mathcal{C}}_2$, *and*

$$\gamma_2 = \frac{\overline{\mathcal{C}}_3^2 - \overline{\mathcal{C}}_4 k + k^3}{k(\overline{\mathcal{C}}_3 + k - k^2)}. \tag{2}$$

(*v*) $p_0 + p_1 + \cdots + p_d = H$, *where* H *is the Hoffman polynomial [20].*

(*vi*) *For every* $i = 0, \ldots, d$, *(any multiple of) the sum polynomial* $q_i = p_0 + \cdots + p_i$ *maximizes the quotient* $r(\theta_0)/\|r\|_\Gamma$ *among the polynomials* $r \in \mathbb{R}_i[x]$ *(notice that* $q_i(\theta_0)^2/\|q_i\|_\Gamma^2 = q_i(\theta_0)$*), and*

$$(1 =) q_0(\theta_0) < q_1(\theta_0) < \cdots < q_d(\theta_0) (= H(\theta_0) = n).$$

A graph G with diameter D is called *m-partially distance-regular*, for some $m = 0, \ldots, D$, if its predistance polynomials satisfy $p_i(\boldsymbol{A}) = \boldsymbol{A}_i$ for every $i \leq m$. In particular, every m-partially distance-regular with $m \geq 1$ must be regular (see Abiad, Van Dam, and Fiol [1]). As an alternative characterization, a graph G is m-partially distance-regular when the intersection numbers c_i $(i \leq m)$, a_i $(i \leq m-1)$, b_i $(i \leq m-1)$ are well-defined. In this case, these intersection numbers are equal to the corresponding preintersection numbers γ_i $(i \leq m)$, α_i $(i \leq m-1)$, β_i $(i \leq m-1)$, and also k_i is well-defined and equal to $p_i(\theta_0)$ for $i \leq m$. We refer to Dalfó, Van Dam, Fiol, Garriga, and Gorissen [7] for more background.

Then, with this definition, a graph Γ with diameter $D = d$ is distance-regular if and only it is d-partially distance-regular. In fact, in this case we have the following strongest proposition, which is a combination of results in Fiol, Garriga and Yebra [18], and Dalfó, Van Dam, Fiol, Garriga and Gorissen [7].

Proposition 2. *A regular graph* Γ *with* $d+1$ *different eigenvalues is distance-regular if and only if there exists a polynomial* p *of degree* d *such that* $p(\boldsymbol{A}) = \boldsymbol{A}_d$, *in which case* $p = p_d$. □

Lemma 3. *Let* Γ *be a regular graph with diameter* D, *and let* $m \leq D$ *be a positive integer. Let* $\overline{n_i} = \frac{1}{n}\sum_{u \in V} n_i(u)$ *be the average number of vertices at distance at most* $i \leq D$ *from every vertex in* Γ. *Then, for any nonzero polynomial* $r \in \mathbb{R}_i[x]$ *we have*

$$\frac{r(\theta_0)^2}{\|r\|_\Gamma^2} \leq \overline{n_i}, \tag{3}$$

with equality if and only if r *is a multiple of* $q_i = p_0 + \cdots + p_i$, *and* $q_i(\boldsymbol{A}) = \boldsymbol{A}_0 + \cdots + \boldsymbol{A}_i$.

Proof. Let $\boldsymbol{S}_i = \boldsymbol{A}_0 + \cdots + \boldsymbol{A}_i$. As $\deg r \leq i$, we have $\langle r(\boldsymbol{A}), \boldsymbol{J}\rangle = \langle r(\boldsymbol{A}), \boldsymbol{S}_i\rangle$. But $\langle r(\boldsymbol{A}), \boldsymbol{J}\rangle = \langle r, H\rangle_\Gamma = r(\theta_0)$. Then the Cauchy-Schwarz inequality gives

$$r^2(\theta_0) \leq \|r(\boldsymbol{A})\|^2 \|\boldsymbol{S}_i\|^2 = \|r\|_\Gamma^2 \overline{n_i},$$

whence (3) follows. Besides, in case of equality we have that r is multiple of q_i, by Lemma 1(vi), with $q_i(\theta_0) = \overline{n_i}$. Therefore, $q_i(\boldsymbol{A}) = \alpha \boldsymbol{S}_i$ for some nonzero constant α and taking norms we conclude that $\alpha = 1$. □

In fact, as it was shown in Fiol [14], the above result still holds if we change the arithmetic mean of the numbers $n_i(u)$, $u \in V$, by its harmonic mean.

As a consequence of Lemma 3 and Proposition 2, we have the following generalization of the spectral excess theorem, due to Fiol and Garriga [17] (for short proofs, see Van Dam [8], and Fiol, Gago and Garriga [16]):

Theorem 4. *Let Γ be a regular graph with $d+1$ distinct eigenvalues and diameter $D = d$. Let $m \leq D$ be a positive integer.*

- (i) *If Γ is $(m-1)$-partially distance-regular for some $m < d$, and $q_m(\theta_0) = \overline{n_m}$, then Γ is m-partially distance-regular.*
- (ii) *If $q_{d-1}(\theta_0) = \overline{n_{d-1}}$, then Γ is distance-regular.*

2 The case of distance-regular graphs

Here we study the case when Γ is a distance-regular graph with diameter three or four. In fact, in the first case everything is basically known (see [2]), although only a combinatorial characterization was provided, whereas we think that the spectral characterization is also important. Indeed, Brouwer [2] proved the following (see also Proposition 4.2.17(i) in [3]):

Proposition 5. *[2] Let Γ be a distance-regular graph with degree k and diameter $d = 3$. Then,*

- (i) *Γ_2 is strongly regular $\iff$ $c_3(a_3 + a_2 - a_1) = b_1 a_2$.*
- (ii) *$\Gamma_{1,2}$ is strongly regular $\iff$ Γ has eigenvalue -1 $\iff$ $k = b_2 + c_3 - 1$.*

Notice that, in this case, $\Gamma_{1,2}$ is strongly regular if and only if its complement Γ_3 is. As commented in the Introduction, the last case was studied for general diameter by Brouwer and Fiol [4] and Fiol [15].

Proposition 6. *Let Γ be a distance-regular graph with diameter $D = d = 3$, and eigenvalues $\theta_0 (= k) > \theta_1 > \theta_2 > \theta_3$.*

- (i) *The distance-2 graph Γ_2 is strongly regular if and only if $a_2 - c_3$ is an eigenvalue of Γ.*

(*ii*) *The distance 1-or-2 graph* $\Gamma_{1,2}$ *is strongly regular if and only if* $a_3 - b_2$ *is an eigenvalue of* Γ.

Proof. We only prove (*i*), as the proof of (*ii*) is similar. As Γ_2 has adjacency matrix $\boldsymbol{A}_2 = p_2(\boldsymbol{A})$, where $p_2(x) = \frac{1}{c_2}(x^2 - a_1 x - k)$, it has eigenvalues $\frac{1}{c_2}(\theta^2 - a_1\theta - k)$, where θ is an eigenvalue of Γ. Assume $\eta^2 - a_1\eta - k = \theta^2 - a_1\theta - k$. This implies $\theta = \eta$ or $\theta + \eta = a_1$. Let τ be the third non-trivial eigenvalue of Γ. Then $k + \theta + \eta + \tau = a_1 + a_2 + a_3$ and the result follows. The other direction is trivial to see. □

For diameter $D = d = 4$ only the case of the distance 1-or-2 graph is known (see Proposition 4.2.18 in [3]). In the following result, we give an equivalent characterization of this case and, moreover, we study the case of the distance 2-graph which, as far as we know, it is new. The proof is as in Proposition 6.

Proposition 7. *Let* Γ *be a distance-regular graph with diameter four and eigenvalues* $\theta_0(= k) > \theta_1 > \theta_2 > \theta_3 > \theta_4$.

(*i*) *The distance-2 graph* Γ_2 *is strongly regular if and only if* $\theta_1 + \theta_4 = a_1 = \theta_2 + \theta_3$.

(*ii*) *The distance 1-or-2 graph* Γ_2 *is strongly regular if and only if* $\theta_1 + \theta_4 = a_1 - c_2 = \theta_2 + \theta_3$.

An example of distance-regular graph satisfying the conditions of Proposition 7 is the Hamming graph $H(4,3)$ (see Example 2 in the next section). Another example would be the (possible) graph corresponding to the feasible array $\{39, 32, 20, 2; 1, 4, 16, 30\}$ (see [3, p. 420]). If it exists, this would be a graph with $n = 768$ vertices and spectrum $39^1, 15^{52}, 7^{117}, -1^{468}, -9^{130}$. In this case, its distance-2 graph would have spectrum 312^1, 24^{182}, -8^{585}.

3 The case of regular graphs

Now we want to conclude the same result as above but only requiring that the graph Γ is regular. In this case, we use the predistance polynomials and preintersection numbers. Notice that now $p_i(\boldsymbol{A})$ is not necessarily the distance-i matrix $\boldsymbol{A}_i$ (usually not even a 0-1 matrix). However, as above, we consider that $p_2(\boldsymbol{A})$ has only three distinct eigenvalues.

3.1 The case of diameter three

We begin with the case of $d = 3$ (that is, assuming that Γ has four distinct eigenvalues).

Theorem 8. *Let Γ be a regular graph with degree k, n vertices, spectrum $\operatorname{sp}\Gamma = \{\theta_0, \theta_1^{m_1}, \theta_2^{m_2}, \theta_3^{m_3}\}$, where $\theta_0(=k) > \theta_1 > \theta_2 > \theta_3$, and preintersection number γ_2 given by (2). Let $\overline{k_3} = \frac{1}{n}\sum_{u\in V} k_3(u)$ be the average number of vertices at distance 3 from every vertex in Γ. Consider the polynomials*

$$s_1(x) = x^2 - (\theta_1 + \theta_3 - \gamma_2)x + \gamma_2 + \theta_2(\theta_1 - \theta_2 + \theta_3), \tag{4}$$

$$s_2(x) = x^2 - (\theta_1 + \theta_2 - \gamma_2)x + \gamma_2 + \theta_3(\theta_1 - \theta_3 + \theta_2). \tag{5}$$

Then,

$$\overline{k_3} \le \frac{n\sum_{i=1}^{3} m_i(s_j(\theta_i) - \tau_j)^2}{\sum_{i=0}^{3} m_i(s_j(\theta_i) - \tau_j)^2}, \quad j = 1, 2, \tag{6}$$

where

$$\tau_j = \frac{s_j(\theta_0)\sum_{i=1}^{3} m_i s_j(\theta_i) - \sum_{i=1}^{3} m_i s_j(\theta_i)^2}{s_j(\theta_0)(n-1) - \sum_{i=1}^{3} m_i s_j(\theta_i)}, \quad j = 1, 2. \tag{7}$$

Equality in (6) holds for some $j \in \{1, 2\}$ if and only if Γ is a distance-regular graph and its distance-2 graph Γ_2 is strongly-regular, with eigenvalues

$$\lambda_0 = n - \overline{k_3} - \theta_0 - 1, \quad \lambda_1 = ((\theta_1 - \theta_2)(\theta_2 - \theta_3) - \tau_1)/\gamma_2, \text{ and } \lambda_2 = -\tau_1/\gamma_2,$$

or

$$\lambda_0 = n - \overline{k_3} - \theta_0 - 1, \quad \lambda_1 = -\tau_2/\gamma_2, \text{ and } \lambda_2 = ((\theta_1 - \theta_3)(\theta_3 - \theta_2) - \tau_2)/\gamma_2,$$

where $k_3 = \overline{k_3}$ is the constant value of the number of vertices at distance 3 from any vertex in Γ.

Proof. Taking into account that the eigenvalues of $p_2(x)$ interlace those of Γ, the only possible cases are:

1. $p_2(\theta_1) = p_2(\theta_3) = \sigma_1/\gamma_2$ and $p_2(\theta_2) = -\tau_1/\gamma_2$,
2. $p_2(\theta_1) = p_2(\theta_2) = \sigma_2/\gamma_2$ and $p_2(\theta_3) = -\tau_2/\gamma_2$,

where σ_j and τ_j, for $j = 1, 2$, are constants. We only prove the first case, as the other is similar. The clue is to apply Lemma 3 with a polynomial $r \in \mathbb{R}_2[x]$ having the desired properties of (any multiple of) q_2. To this end, let us assume that $p_2(\theta_1) = p_2(\theta_3) = \sigma_1/\gamma_2$, and $p_2(\theta_2) = -\tau_1/\gamma_2$ where σ_1 and τ_1 are constants. Thus, if we consider a generic monic polynomial $r(x) = x^2 + \alpha x + \beta = \gamma_2 q_2(x)$, where $q_2(x) = p_2(x) + x + 1$, we must have

$$\begin{aligned} r(\theta_1) &= \theta_1^2 + \alpha\theta_1 + \beta = \sigma_1 + \gamma_2\theta_1 + \gamma_2, \\ r(\theta_2) &= \theta_2^2 + \alpha\theta_2 + \beta = -\tau_1 + \gamma_2\theta_2 + \gamma_2, \\ r(\theta_3) &= \theta_3^2 + \alpha\theta_3 + \beta = \sigma_1 + \gamma_2\theta_3 + \gamma_2. \end{aligned}$$

From the first and last equation we get $\alpha = \gamma_2 - \theta_1 - \theta_3$ and, hence, the second equation yields $\beta = \gamma_2 + \theta_2(\theta_1 - \theta_2 + \theta_3) - \tau_1$. Then, we must take $r(x) = s_1(x) - \tau_1$, where $s_1(x)$ is as in (4), and (3) yields

$$\Phi(\tau_1) = \frac{r(\theta_0)^2}{\|r\|_\Gamma^2} = \frac{n(s(\theta_0) - \tau_1)^2}{\sum_{i=0}^3 m_i(s(\theta_i) - \tau_1)^2} \leq \overline{s_2} = n - \overline{k_3}. \tag{8}$$

Now, to have the best result in (8), and, since we are mostly interested in the case of equality, we find the maximum of the function Φ, which is attained at τ_1 given by (14). Then, as $\overline{s_2} = n - \overline{k_3}$, the claimed inequality follows. Moreover, in case of equality, we know, by Theorem 4, that Γ is distance-regular with $r(x) = \gamma q_{d-1}(x)$ for some constant γ, which it is $\gamma = \gamma_2$. Then, we get

$$P_{22} = p_2(\theta_2) = -\tau_1/\gamma_2, \text{ and } P_{i2} = p_2(\theta_i) = \sigma_1/\gamma_2 = ((\theta_1-\theta_2)(\theta_2-\theta_3)-\tau_1)/\gamma_2 \text{ for } i = 1, 3.$$

To prove the converse, we only need to carry out a simple computation. Indeed, assume that Γ is a distance-regular graph, with k_i being the vertices at distance $i = 1, \ldots, 3$ from any vertex ($k_1 = k$), and $p_2(\theta_1) = p_d(\theta_3)$. Then, the same reasoning as in Proposition 6 gives $a_1 = \theta_1 + \theta_3$. Then, from $kb_1 = c_2k_2 = c_2(n - k_3 - k - 1)$ and $a_1 + b_1 + 1 = k$, we get that $c_2 = \frac{k(k-1-\theta_1-\theta_3)}{n-k_3-k-1}$. Thus, by putting $\gamma_2 = c_2$ in $s_1(x)$ of (4) to compute τ_1 in (7), the inequality (6) becomes an equality (since $\overline{k_3} = k_3$). □

The following result gives similar conditions for Γ to be distance-regular with the distance 1-or 2 graph $\Gamma_{1,2}$ being strongly regular.

Theorem 9. *Let Γ be a regular graph with degree k, n vertices, spectrum $\operatorname{sp}\Gamma = \{\theta_0, \theta_1^{m_1}, \theta_2^{m_2}, \theta_3^{m_3}\}$, where $\theta_0(= k) > \theta_1 > \theta_2 > \theta_3$, and preintersection number γ_2. Let $\overline{k_3} = \frac{1}{n}\sum_{u\in V} k_3(u)$ be the average number of vertices at distance 3 from every vertex in Γ. Consider the polynomials*

$$s_1(x) = x^2 - (\theta_1 + \theta_3)x + \gamma_2 + \theta_2(\theta_1 - \theta_2 + \theta_3), \tag{9}$$

$$s_2(x) = x^2 - (\theta_1 + \theta_2)x + \gamma_2 + \theta_3(\theta_1 - \theta_3 + \theta_2). \tag{10}$$

Then,

$$\overline{k_3} \leq \frac{n\sum_{i=1}^3 m_i(s_j(\theta_i) - \tau_j)^2}{\sum_{i=0}^3 m_i(s_j(\theta_i) - \tau_j)^2}, \quad j = 1, 2, \tag{11}$$

where

$$\tau_j = \frac{s_j(\theta_0)\sum_{i=1}^3 m_i s_j(\theta_i) - \sum_{i=1}^3 m_i s_j(\theta_i)^2}{s_j(\theta_0)(n-1) - \sum_{i=1}^3 m_i s_j(\theta_i)}, \quad j = 1, 2. \tag{12}$$

Equality in (11) *holds with some $j \in \{1, 2\}$ if and only if Γ is a distance-regular graph and its distance 1-or-2 graph $\Gamma_{1,2}$ is strongly-regular,*

Example 1. The Odd graph O(4) with 7 points, has $n = \binom{7}{3} = 35 = 1+4+12+18$ vertices, diameter $d = 3$, intersection array $\{4, 3, 3; 1, 1, 2\}$, and spectrum $4^1, 2^{14}, -1^{14}, -3^6$. Then, the functions $\Phi(\tau_j)$ in (6) with $j = 1, 2$ have maximum values at $\tau_1 = 18/5$ and $\tau_2 = -8$, respectively, and their values are $\Phi(18/5) = 138/7$ and $\Phi(-8) = 22$. Then, since both numbers are greater than $k_3 = 18$, its distance-2 graph Γ_2 is not strongly regular.

On the other hand the function $\Phi(\tau_1)$ in (11) has maximum value at $\tau_1 = 4$, and $\Phi(4) = 18 = k_3$. Hence, its distance 1-or-2 graph $\Gamma_{1,2}$ (and, hence, also Γ_3) is strongly regular with $p_1(x) + p_2(x) = x^2 + x - 4$, and spectrum $16^1, 2^{20}, -4^{14}$.

3.2 The case of diameter four

The following result deals with the case of $d = 4$. As in the case of Theorem 9, we omit this proof as goes along the same lines of reasoning as in Theorem 8.

Theorem 10. *Let Γ be a regular graph with degree k, n vertices, spectrum $\operatorname{sp}\Gamma = \{\theta_0, \theta_1^{m_1}, \theta_2^{m_2}, \theta_3^{m_3}, \theta_4^{m_4}\}$, where $\theta_0(= k) > \theta_1 > \theta_2 > \theta_3 > \theta_4$, such that $\theta_1 + \theta_4 = \theta_2 + \theta_3$, and preintersection number γ_2. Let $\overline{n_2} = \frac{1}{n}\sum_{u\in V} n_2(u)$ be the average number $n_2(u) = |N_2(u)|$ of vertices at distance at most 2 from every vertex u in Γ. Consider the polynomials*

$$s_1(x) = x^2 - (\theta_2 + \theta_3 - \gamma_2)x + \theta_2\theta_3,$$
$$s_2(x) = x^2 - (\theta_2 + \theta_3)x + \gamma_2 - \theta_2\theta_3.$$

Then,

$$\overline{n_2} \geq \Phi(\tau_j) = \frac{n(s_j(\theta_0) - \tau_j)^2}{\sum_{i=0}^4 m_i(s_j(\theta_i) - \tau_j)^2}, \quad j = 1, 2, \tag{13}$$

where

$$\tau_j = \frac{s_j(\theta_0)\sum_{i=1}^4 m_i s_j(\theta_i) - \sum_{i=1}^4 m_i s_j(\theta_i)^2}{s_j(\theta_0)(n-1) - \sum_{i=1}^4 m_i s_j(\theta_i)}, \quad j = 1, 2. \tag{14}$$

Equality in (13) *holds with $j = 1$ or $j = 2$ if and only if Γ is a 2-partially distance-regular graph and its distance-2 or distance 1-or-2 graph, respectively, is strongly-regular.*

Example 2. The Hamming graph $H(4, 3)$, with $n = 3^4 = 81$ vertices and diameter $d = 4$, has intersection array $\{8, 6, 4, 2; 1, 2, 3, 4\}$, so that $k_4 = 16$, and spectrum $8^1, 5^8, 2^{24}, -1^{32}, -4^{16}$. Then, the function $\Phi(\tau_j)$ in (13) with $j = 1$ has a maximum at $\tau_1 = 4$, and its value is $\Phi(4) = 33 = s_2$. Then, $P_{14} = P_{44}$ and $P_{24} = P_{34}$. Indeed, its distance-2 polynomial is $p_2(x) = \frac{1}{2}(x^2 - x - 8)$ with values $p_4(8) = 24$, $p_4(5) = 6$, $p_4(2) = -3$, $p_4(-1) = -3$, and $p_4(-4) = 6$. Hence, the distance-2 graph Γ_2 is strongly regular with spectrum $24^1, 6^{24}, -3^{56}$.

Example 3. The Odd graph O(5) with 9 points, has $n = \binom{9}{4} = 126 = 1 + 5 + 20 + 40 + 60$ vertices, diameter $d = 4$, intersection array $\{5, 4, 4, 3; 1, 1, 2, 2\}$, and spectrum $5^1, 3^{27}, 1^{42}, -2^{48}, -4^8$. Then, the function $\Phi(\tau_j)$ in (13) with $j = 2$ has a maximum at $\tau_2 = 3$, and its value is $\Phi(4) = 26 = s_2$. Then, its distance 1-or-2 polynomial is

$p_{1,2}(x) = p_1(x) + p_2(x) = x^2 + x - 5$ with values $p_{1,2}(5) = 25$, $p_{1,2}(3) = 7$, $p_{1,2}(1) = -3$, $p_{1,2}(-2) = -3$, and $p_{1,2}(-4) = 7$. Hence, the distance 1-or-2 graph $\Gamma_{1,2}$ is strongly regular with spectrum $25^1, 7^{35}, -3^{90}$.

Acknowledgments. This research is supported by the *Ministerio de Ciencia e Innovación*, Spain, and the *European Regional Development Fund* under project MTM2014-60127-P, and the *Catalan Research Council* under project 2014SGR1147 (C. D. and M.A. F.). Also, this work is partially supported by the *National Natural Science Foundation of China* under project No. 11471009, and the *Chinese Academy of Sciences* under its '100 talent' program (J. K.).

References

[1] A. Abiad, E.R. van Dam, and M.A. Fiol, Some spectral and quasi-spectral characterizations of distance-regular graphs, *J. Combin. Theory, Ser. A* **143** (2016), 1–18.

[2] A.E. Brouwer, Distance regular graphs of diameter 3 and strongly regular graphs, *Discrete Math.* **49** (1984) 101–103.

[3] A.E. Brouwer, A.M. Cohen, and A. Neumaier, *Distance-Regular Graphs*, Springer-Verlag, Berlin-New York, 1989.

[4] A.E. Brouwer and M.A. Fiol, Distance-regular graphs where the distance-d graph has fewer distinct eigenvalues, *Linear Algebra Appl.* **480** (2015), 115–126.

[5] A.E. Brouwer and W.H. Haemers, *Spectra of Graphs*, Springer, 2012; available online at `http://homepages.cwi.nl/~aeb/math/ipm/`.

[6] M. Cámara, J. Fàbrega, M.A. Fiol, and E. Garriga, Some families of orthogonal polynomials of a discrete variable and their applications to graphs and codes, *Electron. J. Combin.* **16(1)** (2009), #R83.

[7] C. Dalfó, E.R. van Dam, M.A. Fiol, E. Garriga, and B.L. Gorissen, On almost distance-regular graphs, *J. Combin. Theory, Ser. A* **118** (2011), 1094–1113.

[8] E.R. van Dam, The spectral excess theorem for distance-regular graphs: a global (over)view, *Electron. J. Combin.* **15(1)** (2008), #R129.

[9] E.R. van Dam, J.H. Koolen, and H. Tanaka, Distance-regular graphs, *Electron. J. Combin.* (2016) #DS22.

[10] V. Diego, J. Fàbrega, and M.A. Fiol, Equivalent characterizations of the spectra of graphs and applications to measures of distance-regularity, submitted (2016).

[11] M.A. Fiol, An eigenvalue characterization of antipodal distance-regular graphs, *Electron. J. Combin.* **4** (1997), #R30.

[12] M.A. Fiol, A quasi-spectral characterization of strongly distance-regular graphs, *Electron. J. Combin.* **7** (2000), #R51.

[13] M.A. Fiol, Some spectral characterization of strongly distance-regular graphs, *Combin. Probab. Comput.* **10** (2001), no. 2, 127–135.

[14] M.A. Fiol, Algebraic characterizations of distance-regular graphs, *Discrete Math.* **246** (2002), 111–129.

[15] M.A. Fiol, The spectral excess theorem for distance-regular graphs having distance-d graph with fewer distinct eigenvalues, *J. Algebraic Combin.* **43** (2016), no. 4, 827–836.

[16] M.A. Fiol, S. Gago, and E. Garriga, A simple proof of the spectral excess theorem for distance-regular graphs, *Linear Algebra Appl.* **432** (2010), 2418–2422.

[17] M.A. Fiol and E. Garriga, From local adjacency polynomials to locally pseudo-distance-regular graphs, *J. Combin. Theory Ser. B* **71** (1997), 162–183.

[18] M.A. Fiol, E. Garriga, and J.L.A. Yebra, Locally pseudo-distance-regular graphs, *J. Combin. Theory Ser. B* **68** (1996), 179–205.

[19] C.D. Godsil, *Algebraic Combinatorics*, Chapman and Hall, NewYork, 1993.

[20] A.J. Hoffman, On the polynomial of a graph, *Amer. Math. Monthly* **70** (1963), 30–36.

[21] G. Szegö, *Orthogonal Polynomials*, 4th edition, American Mathematical Society, Providence, R.I., 1975.